

WEIGHT INVARIANCE OF SOBOLEV-*BMO* SPACES AND WEIGHTED HARDY SPACE ESTIMATES

JAROD HART(UNIVERSITY OF KANSAS)

ABSTRACT. In this talk, we discuss some weight invariant type properties for Sobolev-*BMO* spaces, and use the properties to prove weighted Hardy space estimates for Calderón-Zygmund operators. Roughly speaking, Sobolev-*BMO* spaces $I_s(BMO)$ for $s \geq 0$ are made up of s order anti-derivatives of *BMO* functions, where *BMO* is the John-Nirenberg space of Bounded Mean Oscillation. The weighted counterparts of these spaces $I_s(BMO_w)$ are defined by replacing the Lebesgue measure dx in the Sobolev-*BMO* norms with a Muckenhoupt weighted measure $w(x)dx$. These spaces are very resilient to weight perturbations, which we show by proving $I_s(BMO) = I_s(BMO_w)$ for any Muckenhoupt weight $w \in A_\infty$. We use this weight invariant result to prove that a Calderón-Zygmund operator T , satisfying appropriate cancellation conditions, is bounded on the weighted Hardy space H_w^p when $p_0 < p < \infty$ and $w \in A_{p/p_0}$; here $0 < p_0 < 1$ depends on the operator T . Interestingly, these estimates do not collapse to the known Lebesgue space theory for Calderón-Zygmund operators when $1 < p < \infty$. In fact, it is possible for a Calderón-Zygmund operator T to be bounded on H_w^p for $w \in A_q$ when $1 < p < q < \infty$, in which case T is not necessarily bounded on L_w^p . The work presented in this talk is a joint work with Lucas Oliveira.